

Date:

To: Parents/Guardians:

Re: 2024-2025 Food Allergy & Anaphylaxis Emergency Care Plan

Please download, review, and sign the FARE (Food Allergy & Anaphylaxis Emergency Care Plan) form at <http://www.foodallergy.org/file/emergency-care-plan.pdf>. Please complete the entire form, obtain required signatures, and return to your child's school.

The FARE form addresses:

- Severe Symptoms
- Mild Symptoms
- Medication/Doses
- Directions – Epipen Auto Injector
- Directions – Adrenaclick
- Directions – AUVI-Q

In addition, please sign and return this memo along with the FARE form (which requires parent and physician signatures).

As per parent/guardian of the student listed below, I understand that if the procedures as specified in N.J.S.A. 18A:40-12.6 are followed, the district or non public school shall have no liability as a result of any injury arising from the administration of the epinephrine via a pre-filled auto-injector mechanism to the pupil and that the parents or guardians shall indemnify and hold harmless the district, non public school, and its employees or agents against any claims arising out of the administration of the epinephrine via a pre-filled auto-injector mechanism to the pupil.

Student's Name: _____ School: _____

Physician Signature: _____ Date _____ Phone: _____

Parent/Guardian Signature: _____ Date _____ Phone: _____

Thank you

Name: _____ D.O.B.: _____

Allergic to: _____

 Weight: _____ lbs. Asthma: ☐ Yes (higher risk for a severe reaction) ☐ No

 PLACE
PICTURE
HERE

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.
Extremely reactive to the following allergens: _____

THEREFORE:

- ☐ If checked, give epinephrine immediately if the allergen was **LIKELY** eaten, for ANY symptoms.
- ☐ If checked, give epinephrine immediately if the allergen was **DEFINITELY** eaten, even if no symptoms are apparent.

FOR ANY OF THE FOLLOWING: SEVERE SYMPTOMS



LUNG

Shortness of breath, wheezing, repetitive cough



HEART

Pale or bluish skin, faintness, weak pulse, dizziness



THROAT

Tight or hoarse throat, trouble breathing or swallowing



MOUTH

Significant swelling of the tongue or lips



SKIN

Many hives over body, widespread redness



GUT

Repetitive vomiting, severe diarrhea



OTHER

Feeling something bad is about to happen, anxiety, confusion

 OR A
COMBINATION
of symptoms
from different
body areas.

- 1. INJECT EPINEPHRINE IMMEDIATELY.**
- 2. Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
 - Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
 - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
 - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
 - Alert emergency contacts.
 - Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS



NOSE

Itchy or runny nose, sneezing



MOUTH

Itchy mouth



SKIN

A few hives, mild itch



GUT

Mild nausea or discomfort

FOR MILD SYMPTOMS FROM MORE THAN ONE SYSTEM AREA, GIVE EPINEPHRINE.
FOR MILD SYMPTOMS FROM A SINGLE SYSTEM AREA, FOLLOW THE DIRECTIONS BELOW:

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

 Epinephrine Dose: ☐ 0.1 mg IM ☐ 0.15 mg IM ☐ 0.3 mg IM


Antihistamine Brand or Generic: _____

Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____


HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

1. Remove Auvi-Q from the outer case. Pull off red safety guard.
2. Place black end of Auvi-Q against the middle of the outer thigh.
3. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
4. Call 911 and get emergency medical help right away.




HOW TO USE EPIPEN®, EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN AUTO-INJECTOR, MYLAN

1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, remove the blue safety release by pulling straight up.
3. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
4. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.




HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENALCLICK®), USP AUTO-INJECTOR, AMNEAL PHARMACEUTICALS

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip. Grasp the auto-injector in your fist with the red tip pointing downward.
3. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh. Press down hard and hold firmly against the thigh for approximately 10 seconds.
4. Remove and massage the area for 10 seconds. Call 911 and get emergency medical help right away.




HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES

1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, pull off the blue safety release.
3. Place the orange tip against the middle of the outer thigh at a right angle to the thigh.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE SYMJEPITM (EPINEPHRINE INJECTION, USP)

1. When ready to inject, pull off cap to expose needle. Do not put finger on top of the device.
2. Hold SYMJEPITM by finger grips only and slowly insert the needle into the thigh. SYMJEPITM can be injected through clothing if necessary.
3. After needle is in thigh, push the plunger all the way down until it clicks and hold for 2 seconds.
4. Remove the syringe and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.
5. Once the injection has been administered, using one hand with fingers behind the needle slide safety guard over needle.



ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can worsen quickly.

EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: _____
 DOCTOR: _____ PHONE: _____
 PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____ PHONE: _____
 NAME/RELATIONSHIP: _____ PHONE: _____
 NAME/RELATIONSHIP: _____ PHONE: _____

2024-2025 MANDATORY MEDICATION FORM

ALL MEDICATION (prescription and OTC, including Tylenol and Advil) must be accompanied by written permission from BOTH the PARENT and PHYSICIAN.

- **Prescription medication** must be delivered to the nurse by the parent in the original container, labeled with the student's name, medication, dosage and physician's name.
- **OTC medication** must be delivered to school by the parent in the original sealed container and labeled with the student's name.
- **Written permission** of the student's physician and parent/guardian are required, including the student's name, purpose of the medication, the time (or circumstance) at which the medication should be administered, and the length of time for which the medication is prescribed.

Only those medications which are medically necessary during school hours for a student's wellbeing should be sent to school.

NOTE: THE FIRST DOSE OF ANY MEDICATION MAY NOT BE GIVEN AT SCHOOL.

NAME OF STUDENT _____ DOB _____

NAME OF MEDICATION _____

DOSAGE _____

TIME TO BE GIVEN _____

REASON FOR MEDICATION _____

MEDICATION TO BE GIVEN FROM _____ TO _____
DATE DATE

HOW IT IS TAKEN _____

EXAMPLE: BY MOUTH, INHALER, WITH FOOD, CRUSHED, ETC.

ADDITIONAL COMMENTS _____

PARENT SIGNATURE/DATE

PHYSICIAN SIGNATURE/DATE

TELEPHONE NUMBER

TELEPHONE NUMBER

**2024-2025 PHYSICIAN/PARENT CERTIFICATION FOR
STUDENT'S SELF-ADMINISTRATION OF MEDICATION**

CERTIFICATION TO BE COMPLETED BY PHYSICIAN

STUDENT NAME: _____

DIAGNOSIS: _____

NAME OF MEDICATION: _____

DOSAGE: _____

TIME AND CIRCUMSTANCES OF ADMINISTRATION: _____

POSSIBLE SIDE EFFECTS: _____

I certify that _____ has a potentially life threatening illness
(Student)
which requires the use of _____. I further certify that
(Medication)
_____ is capable and has been instructed in the proper method of
(Student)
self-administration of _____
(Medication)

Signature of Physician Date

PHYSICIAN NAME: _____ TELEPHONE #: _____

CERTIFICATION TO BE COMPLETED BY PARENT

I hereby authorize my son/daughter _____ to self-administer (Name
of Medication) _____ in accordance with special guidelines.

I acknowledge that the school shall incur no liability as a result of any injury arising from the self-
administration of medication by (student name) _____.

I shall indemnify and hold harmless the school, its employees and agents against any and all claims arising
out of the self-administration of (medication) _____ by
(student name) _____.

Parent/Guardian Signature Date

SELF-ADMINISTRATION OF MEDICATION IN SCHOOL

Under N.J.S.A. 18A:40-12.3, self-administration of medication by a pupil for asthma or other potentially
life threatening illness is allowed under guidelines established by the school and provided that the statutory
requirements set forth in this form are complied with.

Any permission for the self-administration of medication is effective for this school year only.

N.J.S.A. 18A:40-12.3 PROVIDES THAT THE SCHOOL SHALL INCUR NO LIABILITY AS A
RESULT OF ANY INJURY ARISING FROM THE SELF-ADMINISTRATION OF MEDICATION BY
A STUDENT.

Rev: 4/2015

ADDITIONAL MEDICATIONS

NAME OF STUDENT _____ DOB _____

NAME OF MEDICATION _____

DOSAGE _____

TIME TO BE GIVEN _____

REASON FOR MEDICATION _____

MEDICATION TO BE GIVEN FROM _____ TO _____
DATE DATE

HOW IT IS TAKEN _____
EXAMPLE: BY MOUTH, INHALER, WITH FOOD, CRUSHED, ETC.

ADDITIONAL COMMENTS _____

NAME OF STUDENT _____ DOB _____

NAME OF MEDICATION _____

DOSAGE _____

TIME TO BE GIVEN _____

REASON FOR MEDICATION _____

MEDICATION TO BE GIVEN FROM _____ TO _____
DATE DATE

HOW IT IS TAKEN _____
EXAMPLE: BY MOUTH, INHALER, WITH FOOD, CRUSHED, ETC.

ADDITIONAL COMMENTS _____

PARENT SIGNATURE/DATE

PHYSICIAN SIGNATURE/DATE

TELEPHONE NUMBER

TELEPHONE NUMBER

PERMISSION TO SHARE INFORMATION 2024-2025

As you are aware, everyday each of our students has contact with a variety of staff members; teachers, bus drivers, therapists, assistants, cafeteria workers, and student interns. While your child is in the care of these people, it is important that they are aware of any information that would require special considerations for his or her health and safety.

To comply with privacy laws, I am requesting your permission to share personal information about your child. This would consist of only that information deemed necessary to protect the well-being of your child. Examples of information that could be shared about your child may include; known allergies, special diets or food restriction, and a history of seizures. This may be done in the form of a printed list or verbal contact with those people who will be working closely with your child. If you have specific questions regarding your child, please call me at school. As always, please feel comfortable knowing that any information you do not want shared with anyone will be kept confidential. Thank you.

PLEASE COMPLETE, SIGN BELOW AND RETURN THIS FORM TO YOUR CHILD'S SCHOOL

Child's Name: _____

_____ **Yes, I give permission for personal information about my child to be shared with other staff members if it will protect his/her health and safety.**

_____ **No, I do not give permission for personal information about my child to be shared with other staff members if it will protect his/her health and safety.**

Parent/Guardian Signature

Date